

KARYOTYPE EVOLUTION IN VERTEBRATES

The *Elasmobranchii* (sharks and rays) are one of the two branches of cartilaginous fish. Their usually large genomes and difficulty in preparing chromosomes have hindered studies of genome organization and comparison. Yamaguchi et al. (Genome Res)(1) focused on the sequencing of the zebra shark *Stegostoma tigrinum* (or leopard shark), which displays a smaller genome, and the whale shark, *Rhincodon typus*. The results, together with the cytogenetic preparations, allowed the authors to assemble these two genomes at the chromosomal level. Their achievements represent a starting point for studying karyotype and synteny conservation in a large set of vertebrates. An interesting observation is the differences between elasmobranchs and other vertebrates in the ancestral autosomes that are adopted as sex chromosomes.

Elasmobranchii (sharks and rays) are one of the two branches of cartilaginous fishes. Their usually large genomes and the difficulty in chromosome preparations has hampered studies in genome organization and comparison. Yamaguchi et al. (Genome Res)¹ focused on the sequencing of the zebra shark *Stegostoma tigrinum* (or leopard shark), showing a smaller genome, and on whale shark, *Rhincodon typus*. These achievements along with the cytogenetic preparations, allowed them to assemble these two genomes at chromosomal level. Their results were a starting point to investigate karyotype and synteny conservation among a large set of vertebrates. One interesting observation stemming from their studies is the differences in ancestral autosomes that were adopted as sex chromosomes between elasmobranchs and other vertebrates.

1. <https://genome.cshlp.org/content/33/9/1527.long>